

Short-distance expansion of correlation functions in the charge-symmetric two-dimensional two-component plasma:

Exact results

Gabriel Téllez*

Departamento de Física, Universidad de Los Andes, A.A. 4976, Bogotá, Colombia

Abstract

We determine exactly the short-distance leading behavior of the density correlation functions of a two-dimensional two-component charge-symmetric Coulomb gas composed of point particles, in the whole regime of stability where the coulombic coupling $\beta < 2$. More generally, we compute the short-distance behavior of the effective interaction potential between two external arbitrary charges Q_1 and Q_2 immersed in the plasma, for $\beta|Q_1| < 2$ and $\beta|Q_2| < 2$. We also find the short-distance asymptotics of the density profiles near a single external charge Q immersed in the plasma for $\beta|Q| < 2$.

Keywords: Rigorous results in statistical mechanics, Charged fluids (Theory), Correlation functions.

*Electronic address: gtellez@uniandes.edu.co

I. INTRODUCTION

The two-dimensional charge-symmetric two-component plasma is a two-dimensional Coulomb fluid composed of two species of point particles with charges ± 1 . Each pair of particles (charges q and q') at a distance r from each other interact via the logarithmic Coulomb potential in two dimensions

$$v(r) = -qq' \ln \frac{r}{L} \quad (1.1)$$

where L is an arbitrary length scale. We describe the system with classical (i.e. non quantum) statistical mechanics at a reduced inverse temperature (coulombic coupling) β . Provided $\beta < 2$ the system of point particles is stable against the collapse of particles of opposite sign. We will consider here only the regime $\beta < 2$. To study the regime beyond $\beta = 2$ it is necessary to consider an additional repulsive short-range interaction to avoid the collapse.

The two-component plasma is mappable into the quantum sine-Gordon model [1], which in, two dimensions, is an integrable model [2, 3, 4]. For $\beta < 2$, using results from the sine-Gordon model, the bulk thermodynamic quantities of the two-component plasma have been computed exactly [5]. Also, using recent results for the form factors of the sine-Gordon model [7, 8], the large-distance behavior of the particle correlation functions have been determined exactly [9]. In Ref. [10], the large-distance behavior of the mean-force effective potential between two external charges immersed in the plasma was determined. Also in that same work [10], the large-distance behavior of the density profiles around an external charge in the plasma was computed (see also [11]).

The work presented here is complementary to the ones of Refs. [9, 10, 11]. We are interested here in the exact short-distance expansion of the effective interaction energy between two external charges Q_1 and Q_2 immersed in the plasma. We will extensively study the cases $Q_2 = \pm 1$ and Q_1 arbitrary and determine the short-distance expansion of the density profile of around a single external charge in the plasma. Also we will study the case $Q_1 = 1$ and $Q_2 = \pm 1$ which allows us to determine the short-distance expansion of the density correlation functions. Our main tool will be the operator product expansion of the exponential fields of the sine-Gordon model.

This paper is organized as follows. In Sec. II we will recall some basics facts about the exact solution of the two-component plasma and the sine-Gordon model needed for

our purposes. In Sec. III follows a general discussion on the short-distance expansion of the mean-force potential interaction between two arbitrary external charges Q_1 and Q_2 immersed in the plasma, which is obtained by means of the operator product expansion. In Sec. IV, we will consider the case of charge-like particles $Q_1 Q_2 > 0$ and in Sec. V the case of particles with opposite signs $Q_1 Q_2 < 0$. Both sections IV and V have a similar organization, although some technical details are different. First, we specialize in the case where $|Q_1| = |Q_2|$ (identical or opposite external charges) and we study the mean-force potential between these two external charges. Then, we consider the case where $Q_2 = \pm 1$ is a charge of the plasma and $Q_1 = Q > 0$ is arbitrary and we study the density profile around the charge Q . Then, we consider the case $Q_2 = \pm 1$ and $Q_1 = 1$ and determine the short-distance behavior of the density correlation functions of the plasma. We test our general results valid for arbitrary $\beta < 2$ against the known results for the correlation functions when $\beta = 2$. In the appendices are presented some expansions for particular values of β , Q_1 and Q_2 of the expectation values of exponential fields and of some Coulomb-type integrals (Dotsenko–Fateev integrals [12, 13]) that are needed in the main text.

II. THE TWO-COMPONENT PLASMA AND THE SINE-GORDON MODEL

The statistical mechanics of the two-dimensional two-component plasma are best worked out in the grand canonical ensemble with fugacity ζ (with dimensions $length^{-2}$). Writing down the Boltzmann factor of the system one can notice that only the combination $z = \zeta L^{\beta/2}$ is relevant. We shall still call z the fugacity, although it should be noticed that its dimensions are $length^{\beta/2-2}$.

Doing a Hubbard-Stratonovich transformation, the grand canonical partition function Ξ of the plasma can be written as the partition function of the quantum sine-Gordon model [1]

$$\Xi = \frac{\int \mathcal{D}\phi \exp[-S(z)]}{\int \mathcal{D}\phi \exp[-S(0)]} \quad (2.1)$$

with

$$S(z) = - \int d^2\mathbf{r} \left[\frac{1}{16\pi} \phi \Delta \phi + 2z \cos(b\phi) \right] \quad (2.2)$$

and

$$b^2 = \beta/4. \quad (2.3)$$

Under this mapping, the bulk density and two-body densities of charges $q = \pm 1$ and $q' = \pm 1$ are expressible as [5, 9]

$$n_q^b = z \langle e^{ibq\phi} \rangle \quad (2.4)$$

and

$$n_{qq'}^{(2)}(|\mathbf{r} - \mathbf{r}'|) = z^2 \langle e^{ibq\phi(\mathbf{r})} e^{ibq'\phi(\mathbf{r}')} \rangle \quad (2.5)$$

where the averages are taken with respect to the sine-Gordon action (2.2).

In the sine-Gordon mapping the fugacity z is renormalized by the (diverging) term $e^{\beta v(0)/2}$ [14]. Thus, to make a complete connection between the sine-Gordon model and the statistical mechanics problem of the two-component plasma it is necessary to specify the proper renormalization scheme for the fugacity. In Ref. [4], Zamolodchikov considered the sine-Gordon model as the Gaussian conformal field (with action $S(0)$) perturbed by the $\cos(b\phi)$ operator. Under this scheme, he used the conformal normalization, where the exponential fields $e^{ib\phi}$ are normalized such that

$$\langle e^{ib\phi(\mathbf{x}_1)} \dots e^{ib\phi(\mathbf{x}_n)} e^{-ib\phi(\mathbf{y}_1)} \dots e^{-ib\phi(\mathbf{y}_n)} \rangle_{z \rightarrow 0} = \frac{\prod_{1 \leq j < i \leq n} |\mathbf{x}_i - \mathbf{x}_j|^\beta |\mathbf{y}_i - \mathbf{y}_j|^\beta}{\prod_{i,j=1}^n |\mathbf{x}_i - \mathbf{y}_j|^\beta} \quad (2.6)$$

which is clearly the appropriate normalization for the Coulomb gas system in view of relations (2.4) and (2.5).

The specific dimensionless grand potential of the sine-Gordon model was computed in Ref. [3] in terms of the mass of the soliton of the sine-Gordon model. On the other hand the relationship between the fugacity and the mass of the soliton under the conformal normalization was found in Ref. [4]. With these results the density-fugacity relationship and full thermodynamics of the Coulomb plasma can be obtained [5]. From the fugacity-density relationship from Ref. [5] an explicit formula for the expectation value of the exponential field, which we will need in the following, can be obtained (see also [6])

$$\langle e^{ib\phi} \rangle = 2 \left(\frac{\pi z}{\gamma(\beta/4)} \right)^\xi \left(\frac{\Gamma(\xi/2)}{\Gamma(\frac{1+\xi}{2})} \right)^2 \frac{\tan(\pi\xi/2)}{(4-\beta)\gamma(\beta/4)} \quad (2.7)$$

where we use the notation $\gamma(x) = \Gamma(x)/\Gamma(1-x)$, with $\Gamma(x)$ the Euler gamma function, and $\xi = \beta/(4-\beta)$.

III. THE INTERACTION POTENTIAL BETWEEN TWO EXTERNAL CHARGES AND THE OPERATOR PRODUCT EXPANSION

Let us consider that two external point charges Q_1 and Q_2 are immersed in the plasma and located at the origin and at $\mathbf{r}$ respectively. To avoid the collapse of the external charges with the charges of the plasma of opposite sign, we suppose that $\beta|Q_1| < 2$ and $\beta|Q_2| < 2$. We define the effective interaction mean-force potential $E_{Q_1 Q_2}(r)$ between the charges Q_1 and Q_2 as in [10] by

$$\exp[-\beta E_{Q_1 Q_2}(r)] = \frac{\Xi[Q_1, 0; Q_2, \mathbf{r}]/\Xi}{(\Xi[Q_1]/\Xi)(\Xi[Q_2]/\Xi)} \quad (3.1)$$

where $\Xi[Q_1, 0; Q_2, \mathbf{r}]$ is the grand partition function of the plasma in the presence of the charge Q_1 at the origin and of the charge Q_2 at $\mathbf{r}$ (with the direct interaction $-Q_1 Q_2 \ln r$ between the external charges included in the Hamiltonian) and $\Xi[Q]$ is the grand partition function of the plasma in the presence of a single external charge Q . As shown in Ref. [10], the effective interaction can be expressed in terms of expectation values of exponentials fields of the sine-Gordon model

$$\exp[-\beta E_{Q_1 Q_2}(r)] = \frac{\langle e^{ibQ_1\phi(0)} e^{ibQ_2\phi(\mathbf{r})} \rangle}{\langle e^{ibQ_1\phi} \rangle \langle e^{ibQ_2\phi} \rangle}. \quad (3.2)$$

The denominator in the preceding expression can be computed since an exact expression for the expectation value $\langle e^{ibQ\phi} \rangle$ was conjectured in Ref. [6], a conjecture later supported by several other studies [10, 15, 16, 17],

$$\langle e^{ibQ\phi} \rangle = \left(\frac{\pi z}{\gamma(\beta/4)} \right)^{Q^2 \xi} \exp[I_b(Q)] \quad (3.3)$$

with

$$I_b(Q) = \int_0^\infty \frac{dt}{t} \left[\frac{\sinh^2(2Qb^2t)}{2 \sinh(bt) \sinh(t) \cosh[(1-b^2)t]} - 2Q^2 b^2 e^{-2t} \right]. \quad (3.4)$$

This expression is valid for $\beta|Q| < 2$, otherwise the integral $I_b(Q)$ diverges. Notice that this is the same stability condition necessary to avoid the collapse of an external charge Q with the counterions of the plasma (ions with charge of opposite sign than Q).

On the other hand, the short-distance expansion of $\langle e^{ibQ_1\phi(0)} e^{ibQ_2\phi(\mathbf{r})} \rangle$ can be obtained by means of the operator product expansion [18], which reads [19]

$$e^{ibQ_1\phi(0)} e^{ibQ_2\phi(\mathbf{r})} = \sum_{n=-\infty}^{n=+\infty} C_{Q_1 Q_2}^{n,0}(r) e^{ib(Q_1+Q_2+n)\phi} + \dots \quad (3.5)$$

where the dots denote subdominant contributions from descendant fields (like $(\partial\phi)^2(\bar{\partial}\phi)^2 e^{i(Q_1+Q_2+n)\phi}$ and so on), which will not be needed for our purposes. The coefficients of the operator product expansion have the following form [19]

$$C_{Q_1 Q_2}^{n,0}(r) = z^{|n|} r^{\beta Q_1 Q_2 + n\beta(Q_1+Q_2) + 2|n|(1-\frac{\beta}{4}) + n^2\beta/2} f_{Q_1 Q_2}^{n,0}(z^2 r^{4-\beta}) \quad (3.6)$$

where $f_{Q_1 Q_2}^{n,0}$ admit a power series expansion of the form

$$f_{Q_1 Q_2}^{n,0}(x) = \sum_{k=0}^{\infty} f_k^{n,0}(Q_1, Q_2) x^k. \quad (3.7)$$

The coefficients of the first leading terms of the expansion (3.7) are

$$f_0^{0,0}(Q_1, Q_2) = 1 \quad (3.8)$$

$$f_0^{n,0}(Q_1, Q_2) = j_n(\beta Q_1, \beta Q_2, \beta) \quad \text{for } n > 0 \quad (3.9)$$

$$f_0^{n,0}(Q_1, Q_2) = j_{|n|}(-\beta Q_1, -\beta Q_2, \beta) \quad \text{for } n < 0 \quad (3.10)$$

where

$$j_n(\beta Q_1, \beta Q_2, \beta) = \frac{1}{n!} \int \prod_{k=1}^n d^2 x_k \prod_{k=1}^n |x_k|^{\beta Q_1} |1 - x_k|^{\beta Q_2} \prod_{1 \leq k < j \leq n} |x_k - x_j|^\beta. \quad (3.11)$$

These Coulomb-type integrals have been computed exactly by Dotsenko and Fateev [12, 13]

$$j_n(\beta Q_1, \beta Q_2, \beta) = \pi^n \gamma(\beta/4)^{-n} \prod_{k=1}^n \gamma(k\beta/4) \times \prod_{k=0}^{n-1} \gamma\left(1 + \frac{\beta Q_1}{2} + k\frac{\beta}{4}\right) \gamma\left(1 + \frac{\beta Q_2}{2} + k\frac{\beta}{4}\right) \gamma\left(-1 - \frac{\beta}{2}(Q_1 + Q_2) - (n-1+k)\frac{\beta}{4}\right). \quad (3.12)$$

The coefficient for $n = 0$ of the first subleading term ($k = 1$) in (3.7) reads

$$f_1^{0,0}(Q_1, Q_2) = J(\beta Q_1, \beta Q_2, \beta) \quad (3.13)$$

where

$$J(\beta Q_1, \beta Q_2, \beta) = \int d^2 x d^2 y \frac{|x|^{\beta Q_1} |1 - x|^{\beta Q_2}}{|y|^{\beta Q_1} |1 - y|^{\beta Q_2} |x - y|^\beta} \quad (3.14)$$

which can be expressed in terms of hypergeometric functions ${}_3F_2$ at unity [20, 21]. In Sec. V and in the appendix B we will compute a special case of this integral when $Q_1 = -Q_2$.

Putting all these expressions together, the first terms of the short-distance expansion of the Boltzmann factor of $E_{Q_1 Q_2}(r)$ are

$$\begin{aligned}
\exp[-\beta E_{Q_1 Q_2}(r)] &= \frac{\langle e^{ib(Q_1+Q_2)\phi} \rangle}{\langle e^{ibQ_1\phi} \rangle \langle e^{ibQ_2\phi} \rangle} r^{\beta Q_1 Q_2} \\
&+ \sum_{n=1}^{\infty} \frac{\langle e^{ib(Q_1+Q_2+n)\phi} \rangle}{\langle e^{ibQ_1\phi} \rangle \langle e^{ibQ_2\phi} \rangle} j_n(\beta Q_1, \beta Q_2, \beta) z^n r^{\beta Q_1 Q_2 + n\beta(Q_1+Q_2) + 2n(1-\frac{\beta}{4}) + n^2\beta/2} \{1 + O(z^2 r^{4-\beta})\} \\
&+ \sum_{n=-\infty}^{-1} \frac{\langle e^{ib(Q_1+Q_2+n)\phi} \rangle}{\langle e^{ibQ_1\phi} \rangle \langle e^{ibQ_2\phi} \rangle} j_{|n|}(-\beta Q_1, -\beta Q_2, \beta) z^{|n|} r^{\beta Q_1 Q_2 + n\beta(Q_1+Q_2) + 2|n|(1-\frac{\beta}{4}) + n^2\beta/2} \{1 + O(z^2 r^{4-\beta})\} \\
&+ \frac{\langle e^{ib(Q_1+Q_2)\phi} \rangle}{\langle e^{ibQ_1\phi} \rangle \langle e^{ibQ_2\phi} \rangle} z^2 r^{\beta Q_1 Q_2 + 4 - \beta} J(\beta Q_1, \beta Q_2, \beta) \{1 + O(z^2 r^{4-\beta})\} + O(r^{\beta Q_1 Q_2 + 4}).
\end{aligned} \tag{3.15}$$

The $O(r^{\beta Q_1 Q_2 + 4})$ term in the last line comes from a descendant field contribution which is explicitly computed in Ref. [19], but that will not be needed in the following.

If βQ_1 , βQ_2 and β are small enough the dominant contribution is given by the term $n = 0$ in (3.5)

$$\exp[-\beta E_{Q_1 Q_2}(r)] \underset{r \rightarrow 0}{\sim} \frac{\langle e^{ib(Q_1+Q_2)\phi} \rangle}{\langle e^{ibQ_1\phi} \rangle \langle e^{ibQ_2\phi} \rangle} r^{\beta Q_1 Q_2} \tag{3.16}$$

which is an expected result: the two charges interact at short distances with the bare Coulomb potential $-Q_1 Q_2 \ln r$. Notice however that depending on the values of βQ_1 , βQ_2 or β , some of the following terms in the expansion (3.15) can become dominant over the $n = 0$ term. We will explore this possibility in the following sections.

IV. THE CHARGE-LIKE CASE: $Q_1 > 0$ AND $Q_2 > 0$

Let us consider that Q_1 and Q_2 have the same sign. In the general small- r asymptotics (3.15), the $n = -1$ term will be dominant over the $n = 0$ term when $\beta(Q_1 + Q_2) > 2$, a condition which will take place if $Q_1 + Q_2$ is positive and sufficiently large. On the other hand the term $n = 1$ will be dominant over the $n = 0$ term if $\beta(Q_1 + Q_2) < -2$. As we see, depending on the sign of $Q_1 + Q_2$, the terms of the series for $n < 0$ (if $Q_1 + Q_2 > 0$) or $n > 0$ (if $Q_1 + Q_2 < 0$) will become dominant.

Let us suppose, without loss of generality, that $Q_1 > 0$ and $Q_2 > 0$. The term $-n$ ($n > 0$) will be the dominant term of the series (3.15) if $2 + (n-1)\beta < \beta(Q_1 + Q_2) < 2 + n\beta$. Thus,

the dominant term of the small- r expansion of $\exp[-\beta E_{Q_1 Q_2}(r)]$ is

$$e^{-\beta E_{Q_1 Q_2}(r)} \underset{r \rightarrow 0}{\sim} \begin{cases} \frac{\langle e^{ib(Q_1+Q_2)\phi} \rangle}{\langle e^{ibQ_1\phi} \rangle \langle e^{ibQ_2\phi} \rangle} r^{\beta Q_1 Q_2}, & 0 < \beta(Q_1 + Q_2) < 2 \\ \frac{\langle e^{ib(Q_1+Q_2-n)\phi} \rangle}{\langle e^{ibQ_1\phi} \rangle \langle e^{ibQ_2\phi} \rangle} j_n(-\beta Q_1, -\beta Q_2, \beta) z^n r^{\beta Q_1 Q_2 - n\beta(Q_1+Q_2) + 2n(1-\frac{\beta}{4}) + n^2\beta/2}, & 2 + (n-1)\beta < \beta(Q_1 + Q_2) < 2 + n\beta; \quad n = 1, 2, \dots \end{cases} \quad (4.1)$$

This relation is valid provided we are still in the region of stability of the system, thus $\beta Q_1 < 2$, $\beta Q_2 < 2$ and $\beta < 2$.

There is an interesting relation between the changes of the dominant small- r behavior of the effective potential shown in (4.1) and the corresponding coefficient expressed in terms of the Dotsenko-Fateev integrals (3.11). $j_n(-\beta Q_1, -\beta Q_2, \beta)$ expressed as an integral in equation (3.11) can be interpreted as the partition function of a Coulomb gas with the external charges Q_1 at 0 and Q_2 at 1 and n charges -1 . Thus, the behavior of $E_{Q_1 Q_2}(r)$ when $2 + (n-1)\beta < \beta(Q_1 + Q_2) < 2 + n\beta$ is dominated by the approach of n charges -1 to the positive external charges Q_1 and Q_2 . This confirms and generalizes the analysis of Hansen and Viot [22].

The integral (3.11) defining $j_n(-\beta Q_1, -\beta Q_2, \beta)$ converges at $x_k \rightarrow 0$ if $\beta Q_1 < 2$, at $x_k \rightarrow 1$ if $\beta Q_2 < 2$ and at $|x_k| \rightarrow \infty$ if $\beta(Q_1 + Q_2) > 2 + (n-1)\beta$. On the other hand, $\langle e^{ib(Q_1+Q_2-n)\phi} \rangle$, converges if $\beta(Q_1 + Q_2) < 2 + n\beta$. These are precisely the same conditions for the corresponding term in the small- r expansion (3.15) to become dominant. Thus when the term $-n$ of the general operator product expansion (3.15) becomes dominant, its corresponding coefficient in (4.1) is properly defined.

When the condition $\beta(Q_1 + Q_2) > 2 + (n-1)\beta$ is not satisfied, the integral defining j_n in equation (3.11) is not convergent, but the corresponding term in (3.15) is not the dominant one. To evaluate this subdominant contribution one has to resort to the analytic continuation of j_n provided by the expression (3.12) in terms of Gamma functions.

At $\beta(Q_1 + Q_2) = 2 + (n-1)\beta$, the term $-n$ and the term $-n+1$ of (3.15) are of the same order, in $r^{\beta Q_1 Q_2 - n(n-1)\beta/2}$, and they should be added to obtain the correct leading behavior. The coefficient of the term $-n$ exhibits a pole of first order due to $j_n(-\beta Q_1, -\beta Q_2, \beta)$, but the coefficient of the term $-n+1$ of the expansion (3.15) also exhibits a pole of first order due to $\langle e^{ib(Q_1+Q_2-n+1)\phi} \rangle$. Indeed, if we write $\beta(Q_1 + Q_2) = 2 + (n-1)\beta + \beta\epsilon$, we have [10]

$$\langle e^{ib(Q_1+Q_2-n+1)\phi} \rangle \underset{\epsilon \rightarrow 0}{\sim} -\langle e^{ib(\frac{1}{2b^2}-1)} \rangle \frac{2\pi z}{\beta\epsilon} \quad (4.2)$$

and

$$\begin{aligned}
j_n(-\beta Q_1, -\beta Q_2, \beta) &= \pi j_{n-1}(-\beta Q_1, -\beta Q_2, \beta) \frac{\gamma(\frac{n\beta}{4})\gamma(\frac{\beta}{4} + \frac{\beta\epsilon}{2})}{\gamma(\frac{n\beta}{4} + \frac{\beta\epsilon}{2})} \times \\
&\quad \gamma(1 - \beta Q_1 + \frac{(n-1)\beta}{4})\gamma(\beta Q_1 + \frac{(n-1)\beta}{4} - \frac{\beta\epsilon}{2})\gamma(\frac{\beta\epsilon}{2}) \\
&\underset{\epsilon \rightarrow 0}{\sim} j_{n-1}(-\beta Q_1, -\beta Q_2, \beta) \frac{2\pi}{\beta\epsilon}.
\end{aligned} \tag{4.3}$$

Both coefficients turn out to have the same residue in absolute value but with opposite signs, and thus cancel each other, giving a finite value, and an additional $r^{\beta Q_1 Q_2 - n(n-1)\beta/2} \ln r$ contribution to the small- r expansion. This will be illustrated in several particular cases in the following sections.

A. Effective mean-force potential between two identical external charges

In this section we consider two identical external charges immersed in the plasma $Q_1 = Q_2 = Q > 0$. Using equation (4.1), the first few changes in the small- r asymptotics of the effective potential read

$$e^{-\beta E_{QQ}(r)} \underset{r \rightarrow 0}{\sim} \begin{cases} \frac{\langle e^{2ibQ\phi} \rangle}{\langle e^{ibQ\phi} \rangle^2} r^{\beta Q^2} & 0 < \beta Q < 1 \\ \frac{\langle e^{ib(2Q-1)\phi} \rangle}{\langle e^{ibQ\phi} \rangle^2} j_1(-\beta Q, -\beta Q, \beta) z r^{\beta Q^2 - 2(\beta Q - 1)} & 1 < \beta Q < 1 + \beta/2 \\ \frac{\langle e^{ib(2Q-2)\phi} \rangle}{\langle e^{ibQ\phi} \rangle^2} j_2(-\beta Q, -\beta Q, \beta) z^2 r^{\beta Q^2 - 4(\beta Q - 1) + \beta} & 1 + \frac{\beta}{2} < \beta Q < 1 + \beta; \beta < 1 \\ \frac{\langle e^{ib(2Q-n)\phi} \rangle}{\langle e^{ibQ\phi} \rangle^2} j_n(-\beta Q, -\beta Q, \beta) z^n r^{\beta Q^2 - 2n\beta Q + 2n(1 - \frac{\beta}{4}) + n^2\beta/2} & 1 + \frac{(n-1)\beta}{2} < \beta Q < 1 + \frac{n\beta}{2}; \beta < \frac{2}{n-1}; n = 3, 4, \dots \end{cases} \tag{4.4}$$

Notice that in order to satisfy the relation $\beta Q > 1 + (n-1)\beta/2$ and the stability condition $\beta Q < 2$, the coulombic coupling must be small enough: $\beta < 2/(n-1)$. Thus, as β becomes larger there are few changes in the small- r behavior of the effective potential.

B. Coion density profile around an single external charge $Q > 0$

Let us consider now a single external charge $Q > 0$ at the origin immersed in the plasma. The density of the particles with charge +1 (coions) of the plasma is given by [10, 11]

$$n_+(r) = z \frac{\langle e^{ibQ\phi(0)} e^{ib\phi(\mathbf{r})} \rangle}{\langle e^{ibQ\phi} \rangle} \tag{4.5}$$

or equivalently, using relation (2.4),

$$n_+(r) = n_+^b \frac{\langle e^{ibQ\phi(0)} e^{ib\phi(\mathbf{r})} \rangle}{\langle e^{ibQ\phi} \rangle \langle e^{ib\phi} \rangle} \quad (4.6)$$

Thus, using the expansion (3.15) with $Q_1 = Q$ and $Q_2 = +1$ we can deduce the small- r asymptotics of $n_+(r)$.

To ensure the stability of the system we require that $\beta Q < 2$. This condition reduces the number of changes in the small- r behavior (4.1) to two. Writing explicitly the j_1 coefficient we find that the dominant behavior of the coion density is

$$n_+(r) \underset{r \rightarrow 0}{\sim} \begin{cases} z \frac{\langle e^{ib(Q+1)\phi} \rangle}{\langle e^{ibQ\phi} \rangle} r^{\beta Q}, & 0 < \beta Q < 2 - \beta \\ \pi z^2 \gamma \left(1 - \frac{\beta Q}{2}\right) \gamma \left(1 - \frac{\beta}{2}\right) \gamma \left(-1 + \frac{\beta(Q+1)}{2}\right) r^{2-\beta}, & 2 - \beta < \beta Q < 2 \end{cases} \quad (4.7)$$

Where $\langle e^{ib(Q+1)\phi} \rangle$ and $\langle e^{ibQ\phi} \rangle$ can be computed from equation (3.3). This confirms and completes the analysis of Ref. [11], where the change of behavior at $\beta Q = 2 - \beta$ of the coion density was identified and interpreted as a “precursor” of the counterion condensation, in the sense that for $\beta Q > 2 - \beta$, the effective potential (times β) of a coion and the charge Q behaves at short distances as the bare Coulomb potential with a reduced charge: $-(2 - \beta) \ln r$, instead of $-\beta Q \ln r$. The charge is reduced because of the counterion condensation. With the present work, we have also an explicit expression for the coefficient multiplying $r^{2-\beta}$.

It is also interesting to study the transition at $\beta Q = 2 - \beta$ and at $\beta Q = 2$. At $\beta Q = 2 - \beta$, both terms $r^{\beta Q}$ and $r^{2-\beta}$ shown in equation (4.7) become of the same order and should be added to obtain the correct dominant behavior.

Using the expansions of the expectation values of exponentials field (A2) and (A6) computed in the appendix A we have

$$\begin{aligned} & \frac{\langle e^{ib(Q+1)\phi} \rangle}{\langle e^{ibQ\phi} \rangle} \underset{\beta Q \rightarrow 2-\beta}{=} \frac{2\pi z}{2 - \beta(Q+1)} \times \\ & \left\{ 1 + \frac{2 - \beta(Q+1)}{\beta} \left[R_1(b^2) + I'_b\left(\frac{1}{2b^2} - 1\right) - \frac{2\beta}{4 - \beta} \ln \left(\frac{\pi z}{\gamma(b^2)} \right) \right] + O([\beta(Q+1) - 2]^2) \right\} \end{aligned} \quad (4.8)$$

where $I'_b(Q) = \partial I_b(Q) \partial Q$ and $R_1(b^2)$ are given in equations (A3) and (A7) of the appendix A). On the other hand the $\gamma(-1 + \beta(Q+1)/2)$ from j_1 also exhibit a pole at

$\beta Q = 2 - \beta$, but with opposite residue:

$$\pi z \gamma \left(1 - \frac{\beta Q}{2}\right) \gamma \left(1 - \frac{\beta}{2}\right) \gamma \left(-1 + \frac{\beta(Q+1)}{2}\right) \Big|_{\beta Q \rightarrow 2-\beta} = \frac{2\pi z}{2 - \beta(Q+1)} \left[-1 - \frac{2 - \beta(Q+1)}{2} [2C + \psi(1 - \beta/2) + \psi(\beta/2)] + O([\beta(Q+1) - 2]^2) \right] \quad (4.9)$$

where $\psi(x) = \Gamma'(x)/\Gamma(x)$ is the digamma function and $C = -\psi(1)$ is the Euler constant. Thus adding both contribution gives a finite result. Using (4.8), (4.9) and expanding $r^{\beta Q} = r^{2-\beta}(1 + [\beta(Q+1) - 2] \ln r + O([\beta(Q+1) - 2]^2))$ finally yields, for $\beta Q = 2 - \beta$,

$$n_+(r) \underset{r \rightarrow 0}{\sim} 2\pi z^2 r^{2-\beta} \times \left\{ -\ln \left[\left(\frac{\pi z}{\gamma(b^2)} \right)^{\frac{2}{4-\beta}} r \right] + \frac{1}{\beta} \left[R_1(b^2) + I'_b \left(\frac{1}{2b^2} - 1 \right) \right] - C - \frac{1}{2} [\psi(1 - \beta/2) + \psi(\beta/2)] \right\} \quad (4.10)$$

Notice that the power of z in the logarithm, $z^{2/(4-\beta)}$, is the appropriate one to have a dimensionless argument in the logarithm.

When $\beta Q = 2$, the contribution from the term $n = -2$ of (3.15) in $r^{-\beta Q - \beta + 4}$, is of the same order as the one for $n = -1$, in $r^{2-\beta}$. The coefficients of both terms exhibit a pole with opposite residues when $\beta Q = 2$. A similar mechanism as before applies, adding both contributions gives a finite result. The term $n = -2$ of the expansion of $n_+(r)$ reads

$$z^3 r^{-\beta Q - \beta + 4} \frac{\langle e^{ib(Q-1)\phi} \rangle}{\langle e^{ibQ\phi} \rangle} j_2(-\beta Q, -\beta, \beta) \quad (4.11)$$

with

$$j_2(-\beta Q, -\beta, \beta) = \pi^2 \gamma \left(1 - \frac{\beta}{4}\right)^2 \gamma \left(1 - \frac{\beta Q}{2}\right) \gamma \left(-1 + \frac{\beta Q}{2}\right) \gamma \left(-1 + \frac{\beta Q}{2} + \frac{\beta}{4}\right) \gamma \left(1 - \frac{\beta Q}{2} + \frac{\beta}{4}\right) \quad (4.12)$$

When $\beta Q \rightarrow 2$, we have

$$\frac{\langle e^{ib(Q-1)\phi} \rangle}{\langle e^{ibQ\phi} \rangle} j_2(-\beta Q, -\beta, \beta) \Big|_{\beta Q \rightarrow 2} = \frac{2\pi}{z(\beta Q - 2)} \times \left\{ 1 + \frac{\beta Q - 2}{\beta} \left[R_1(b^2) + I'_b \left(\frac{1}{2b^2} - 1 \right) - 2\xi \ln \frac{\pi z}{\gamma(b^2)} \right] + O((\beta Q - 2)^2) \right\} \quad (4.13)$$

while from the term $n = -1$ we have a contribution from the j_1 coefficient:

$$\gamma \left(1 - \frac{\beta Q}{2}\right) \gamma \left(1 - \frac{\beta}{2}\right) \gamma \left(-1 + \frac{\beta(Q+1)}{2}\right) \Big|_{\beta Q \rightarrow 2} = \frac{2}{2 - \beta Q} - 2C - \psi(1 - \beta/2) - \psi(\beta/2). \quad (4.14)$$

Expanding $r^{-\beta Q - \beta + 4} = r^{2-\beta}[1 + (2 - \beta Q) \ln r + O((2 - \beta Q)^2)]$, and adding all contributions (4.13) and (4.14), we actually find that at $\beta Q = 2$, the small- r behavior of $n_+(r)$ is exactly the same as the one for $\beta Q = 2 - \beta$ given by equation (4.10).

C. Density correlation function $n_{++}^{(2)}(r)$

The two-body density correlation function between particles of the same sign (say positive), can be expressed as

$$n_{++}^{(2)}(r) = z^2 \langle e^{ib\phi(0)} e^{ib\phi(\mathbf{r})} \rangle = (n_+^b)^2 \frac{\langle e^{ib\phi(0)} e^{ib\phi(\mathbf{r})} \rangle}{\langle e^{ib\phi} \rangle^2}. \quad (4.15)$$

Thus, using the results of the last section with $Q = 1$ allow us to find the small- r behavior of the correlation function:

$$n_{++}^{(2)}(\mathbf{r}) \underset{r \rightarrow 0}{\sim} \begin{cases} z^2 \langle e^{2ib\phi} \rangle r^\beta & \beta < 1 \\ 2\pi z^2 n_+^b r \left\{ -\ln \left[\left(\frac{\pi z}{\gamma(1/4)} \right)^{2/3} r \right] + R_1(1/4) + I'_{1/2}(1) + 2 \ln 2 \right\} & \beta = 1 \\ \pi z^2 n_+^b \gamma(1 - \frac{\beta}{2})^2 \gamma(\beta - 1) r^{2-\beta} = \pi z^3 \langle e^{ib\phi} \rangle \gamma(1 - \frac{\beta}{2})^2 \gamma(\beta - 1) r^{2-\beta} & 1 < \beta < 2 \end{cases} \quad (4.16)$$

A numerical estimate of the constant intervening in the case $\beta = 1$ is $R_1(1/4) + I'_{1/2}(1) \simeq -1.73246$.

Our result (4.16) confirms the change of behavior of the correlation function $n_{++}^{(2)}(r)$ at $\beta = 1$, predicted by Hansen and Viot [22]. Furthermore, we obtained an explicit analytical expression for the coefficients multiplying r^β (for $\beta < 1$) and $r^{2-\beta}$ (for $\beta > 1$).

The limit $\beta \rightarrow 2$ cannot be obtained from the results of the previous section because there is an additional divergence due to n_+^b . When $\beta \rightarrow 2$, the term $n = -1$ of the general series (3.15), in $r^{2-\beta}$ is of the same order as the term $n = -2$. This term ($n = -2$) gives a contribution to $n_{++}^{(2)}(r)$ equal to

$$\pi^2 z^4 r^{4-2\beta} \gamma(1 - \frac{\beta}{4})^3 \gamma(1 - \frac{\beta}{2}) \gamma(\frac{\beta}{2} - 1) \gamma(-1 + \frac{3\beta}{4}) = -\frac{4\pi^2 z^4 r^{4-2\beta}}{(2 - \beta)^2} \gamma(1 - \frac{\beta}{4})^3 \gamma(-1 + \frac{3\beta}{4}). \quad (4.17)$$

Notice that the coefficients of $r^{2-\beta}$ in (4.16) and of $r^{4-2\beta}$ in (4.17) exhibit now a pole of second order when $\beta \rightarrow 2$, as opposed to the situation of the previous section, when the pole was of order one. Indeed,

$$\frac{-4\gamma(1 - \frac{\beta}{4})^3 \gamma(-1 + \frac{3\beta}{4})}{(\beta - 2)^2} \underset{\beta \rightarrow 2}{=} -\frac{4}{(\beta - 2)^2} + O(\beta - 2) \quad (4.18)$$

while, using the expansion (A9) of $\langle e^{ib\phi} \rangle$, and expanding the gamma functions,

$$\begin{aligned} \langle e^{ib\phi} \rangle \gamma(1 - \frac{\beta}{2})^2 \gamma(\beta - 1) \underset{\beta \rightarrow 2}{=} \frac{8\pi z}{(\beta - 2)^2} \left[1 + (\beta - 2)[C + \ln(\pi z)] \right. \\ \left. + \frac{(\beta - 2)^2}{2} [C + \ln(\pi z)][1 + C + \ln(\pi z)] \right] + O(\beta - 2) \end{aligned} \quad (4.19)$$

Notice that now the poles do not cancel each other. Expanding $r^{2-\beta} = 1 + (2 - \beta) \ln r + \frac{(2-\beta)^2}{2} (\ln r)^2 + O((\beta - 2)^3)$ and $r^{4-2\beta} = 1 + 2(2 - \beta) \ln r + 2(2 - \beta)^2 (\ln r)^2 + O((\beta - 2)^3)$, and putting together all contributions from (4.18) and (4.19), we finally find

$$\begin{aligned} n_{++}^{(2)}(r) \underset{r \rightarrow 0, \beta \rightarrow 2}{\sim} (2\pi z^2)^2 \left[\frac{1}{(\beta - 2)^2} + \frac{2}{\beta - 2} [C + \ln(\pi z)] \right. \\ \left. - [\ln(\pi z r) + C]^2 + 2[C + \ln(\pi z)]^2 + C + \ln(\pi z) + O(\beta - 2) \right] \end{aligned} \quad (4.20)$$

In the preceding expression we recognize the expansion (A10) of $\langle e^{ib\phi} \rangle^2$ when $\beta \rightarrow 2$. Indeed, we have

$$n_{++}^{(2)}(r) \underset{r \rightarrow 0, \beta \rightarrow 2}{\sim} -(2\pi z^2)^2 (\ln(\pi z r) + C)^2 + z^2 \langle e^{ib\phi} \rangle^2 \quad (4.21)$$

Remembering that $n_+^b = z \langle e^{ib\phi} \rangle$, we recover a known result at $\beta = 2$. The correlation function $n_{++}^{(2)}(r)$ does not have a finite limit when $\beta = 2$, however the truncated correlation function $n_{++}^{(2)T}(r) = n_{++}^{(2)}(r) - (n_+^b)^2$, does have a finite limit at the collapse point $\beta = 2$

$$n_{++}^{(2)T}(r) \underset{r \rightarrow 0}{\sim} -(2\pi z^2)^2 (\ln(\pi z r) + C)^2, \quad \beta = 2 \quad (4.22)$$

Furthermore, the density correlation functions of the two-component plasma at $\beta = 2$ have been computed exactly in Refs. [23, 24]

$$n_{++}^{(2)T}(r) = -(2\pi z^2)^2 (K_0(2\pi z r))^2 \quad (4.23)$$

where K_0 is the modified Bessel function of order 0. Using the well-known expansion of K_0 for small-argument [25] we verify that the small- r asymptotics (4.22) are in complete agreement with the exact expression (4.23).

V. THE OPPOSITE CHARGE CASE: $Q_1 > 0$ AND $Q_2 < 0$

If Q_1 and Q_2 have different signs, say $Q_1 > 0$ and $Q_2 < 0$, then the dominant behavior of $E_{Q_1 Q_2}(r)$ is always given by (3.16), since we always have $\beta|Q_1 + Q_2| < 2$, because in order to

satisfy the stability condition $0 < \beta Q_1 < 2$ and $-2 < \beta Q_2 < 0$. The condition for the term $n = -1$ (or $n = +1$) to become dominant, exposed in Sec. IV, is never satisfied. Therefore, in this section, we will be interested in the next subdominant contribution to $E_{Q_1 Q_2}(r)$ in some particular cases.

A. Effective mean-force potential between two opposite external charges

In this section we are interested in the case $Q_1 = Q > 0$ and $Q_2 = -Q$. Notice that in this case in the general small- r expansion (3.15), the term $n > 0$ and $-n$ are of the same order: $r^{-\beta Q^2 + 2n(1-\beta/4) + n^2\beta/2}$. Furthermore besides the factor $r^{-\beta Q^2}$ which is present for all n , the rest of the power of r is independent of Q . Thus to study the first subdominant contribution we need to take into account both terms $n = 1$ and $n = -1$. Actually, a closer look to the expansion (3.15) shows that in the limit $Q_1 \rightarrow -Q_2$, the coefficients of the expansion diverge. But the coefficient for n have a first order pole with opposite residue than the one for $-n$, thus adding both contributions gives a finite result. Expanding the coefficients for $n = -1$ and $n = 1$ in (3.15) when $Q_1 + Q_2 \rightarrow 0$ and adding both contributions gives

$$\begin{aligned}
e^{-\beta E_{Q,-Q}(r)} \underset{r \rightarrow 0}{=} & \frac{r^{-\beta Q}}{\langle e^{ibQ\phi} \rangle^2} - \pi z r^{-\beta Q+2} \frac{\langle e^{ib\phi} \rangle}{\langle e^{ibQ\phi} \rangle^2} \left(\frac{\beta Q}{2} \right)^2 \times \\
& \left[\frac{4}{\beta} I'_b(1) + 4 \ln \left[\left(\frac{\pi z}{\gamma(b^2)} \right)^{\frac{2}{4-\beta}} r \right] - 4 + 4C + \psi\left(-\frac{\beta Q}{2}\right) + \psi\left(\frac{\beta Q}{2}\right) + \psi\left(1 - \frac{\beta Q}{2}\right) + \psi\left(1 + \frac{\beta Q}{2}\right) \right] \\
& + \frac{z^2}{\langle e^{ibQ\phi} \rangle^2} r^{-\beta Q^2+4-\beta} J(\beta Q, -\beta Q, \beta) + O(r^{-\beta Q^2+4}) \quad (5.1)
\end{aligned}$$

We have also written explicitly the second subdominant contribution of order $r^{-\beta Q+4-\beta}$ which actually does not come from the $n = -2$ and $n = 2$ terms but from the first subdominant contribution to the $n = 0$ term: the last term in equation (3.15). An explicit expression for $J(\beta Q, -\beta Q, \beta)$ is given in the appendix B, formulas (B2-B6) .

The next subdominant contribution $O(r^{-\beta Q^2+4})$, not written here explicitly comes from a contribution of the descendant field $\langle (\partial\phi)^2(\bar{\partial}\phi)^2 \rangle$ [19].

B. Counterion density profile around an single external charge $Q > 0$

The density of negative particles (counterions) of the plasma around a single guest charge $Q > 0$ is given by

$$n_-(r) = z \frac{\langle e^{ibQ\phi(0)} e^{-ib\phi(r)} \rangle}{\langle e^{ibQ\phi} \rangle}. \quad (5.2)$$

The leading term in the small- r expansion is of order $r^{-\beta Q}$, as explained earlier. Using the general expansion (3.15) we find that next subleading contributions comes from the $n = 1$ term if $Q < 1$ and from the $n = -1$ term if $Q > 1$. Thus,

$$n_-(r) \underset{r \rightarrow 0}{=} \begin{cases} z \frac{\langle e^{ib(Q-1)\phi} \rangle}{\langle e^{ibQ\phi} \rangle} r^{-\beta Q} + z^2 r^{2-\beta} j_1(\beta Q, -\beta, \beta) + O(r^{4-\beta-\beta Q}) & Q < 1 \\ z \frac{\langle e^{ib(Q-1)\phi} \rangle}{\langle e^{ibQ\phi} \rangle} r^{-\beta Q} + z^2 \frac{\langle e^{ib(Q-2)\phi} \rangle}{\langle e^{ibQ\phi} \rangle} r^{-2\beta Q + \beta + 2} j_1(-\beta Q, \beta, \beta) + O(r^{4-\beta-\beta Q}, r^{4+3\beta-3\beta Q}) & Q > 1 \end{cases} \quad (5.3)$$

Notice that if $Q = 1$ both subleading contributions presented above are of the same order, in $r^{2-\beta}$, and they should be added as explained in the preceding section. The case $Q = 1$ will be studied in the next section.

The limit $\beta Q \rightarrow 2$ is quite unnatural here. Notice that when $\beta Q \rightarrow 2$, the denominator of the leading term in $r^{-\beta Q}$ diverges. Then this term will disappear when $\beta Q = 2$. On the other hand the next subleading term, from (5.3), case $Q > 1$, has a finite limit when $\beta Q \rightarrow 2$. Indeed, in the numerator $j_1(-\beta Q, \beta, \beta)$ has a pole of order 1 at $\beta = 2$, but so does $\langle e^{ibQ\phi} \rangle$ in the denominator. Thus, we find

$$n_+(r) \underset{r \rightarrow 0}{\sim} z \frac{\langle e^{ib(\frac{1}{2b^2}-2)\phi} \rangle}{\langle e^{ib(\frac{1}{2b^2}-1)\phi} \rangle} r^{\beta-2}, \quad \beta Q = 2. \quad (5.4)$$

We call this limit “unnatural” because the power of r changes in a non-continuous manner from $-\beta Q$ to $\beta - 2$, contrary to the “natural” limits found in Sec. IV. Also the mechanism for finding a finite result is not the same as in Sec. IV, where we added terms of the same order and their poles canceled each other.

Let us recall the fact that at $\beta Q = 2$, strictly speaking we need to add a short-distance regularization to the potential created by the external charge, for example by considering that the external charge is not a point but a small impenetrable disk. With this regularization the denominator in the leading term of (5.3) will not vanish when $\beta Q = 2$ and the density $n_-(r)$ will still behave as $r^{-\beta Q}$, but with a prefactor which is not universal, in the sense that it depends on the regularization procedure (it depends on the radius of the

external disk particle). This is opposed to the situation for $n_+(r)$ that has the universal limit (4.10) when $\beta Q = 2$, independent of the regularization procedure (provided the radius of the external particle is small enough).

C. Density correlation function $n_{+-}^{(2)}(r)$

Putting $Q = 1$ in equation (5.1) allow us to find the small- r expansion of the density correlation function $n_{+-}^{(2)}(r)$ between a positive charge of the plasma and a negative one,

$$n_{+-}^{(2)}(r) \underset{r \rightarrow 0}{=} z^2 r^{-\beta} \left\{ 1 - \pi z r^2 \frac{\beta^2}{4} \langle e^{ib\phi} \rangle \times \right. \\ \left[\frac{4}{\beta} I'_b(1) + 4 \ln \left[\left(\frac{\pi z}{\gamma(b^2)} \right)^{\frac{2}{4-\beta}} r \right] - 4 + 4C + \psi\left(\frac{\beta}{2}\right) + \psi\left(-\frac{\beta}{2}\right) + \psi\left(1 - \frac{\beta}{2}\right) + \psi\left(1 + \frac{\beta}{2}\right) \right] \\ \left. + z^2 r^{4-\beta} J(\beta, -\beta, \beta) + O(r^4) \right\} \quad (5.5)$$

An explicit expression for the coefficient $J(\beta, -\beta, \beta)$ is given in the appendix B, see equations (B7-B9).

As a test, it is interesting to study the limit $\beta \rightarrow 2$ and compare it with the exact expression from Refs. [23, 24]

$$n_{+-}^{(2)T}(r) = (2\pi z^2)(K_1(2\pi z r))^2, \quad \beta = 2 \quad (5.6)$$

where $K_1(x)$ is the modified Bessel of order 1.

Notice that when $\beta \rightarrow 2$ the terms in $r^{2-\beta}$ and $r^{4-2\beta}$ from equation (5.5) become of the same order. Using the expansions (A9) for $\langle e^{ib\phi} \rangle$ and (A12) for $I'_b(1)$ from appendix A, and the expansion (B20) from appendix B for $J(\beta, -\beta, \beta)$ when $\beta \rightarrow 2$, we find

$$n_{+-}^{(2)}(r) \underset{r \rightarrow 0, \beta \rightarrow 2}{=} z^2 r^{-2} + (2\pi z^2)^2 \left\{ \frac{1}{(\beta-2)^2} + \frac{2}{\beta-2} (C + \ln(\pi z)) - \frac{1}{2} + C + \ln(\pi z r) \right. \\ \left. + C + \ln(\pi z) + 2(\ln(\pi z) + C)^2 \right\} + O(r^2) + O(\beta-2). \quad (5.7)$$

Using the expansion (A10) of $\langle e^{ib\phi} \rangle^2$ we can write

$$n_{+-}^{(2)}(r) \underset{r \rightarrow 0, \beta \rightarrow 2}{=} z^2 r^{-2} + (2\pi z^2)^2 \left[-\frac{1}{2} + C + \ln(\pi z r) \right] + z^2 \langle e^{ib\phi} \rangle^2 + O(r^2) + O(\beta-2). \quad (5.8)$$

We recover the well-known fact that although $n_{+-}^{(2)}(r)$ does not have a finite limit when $\beta \rightarrow 2$, the truncated correlation function $n_{+-}^{(2)T}(r) = n_{+-}^{(2)}(r) - n_+^b n_-^b = n_{+-}^{(2)}(r) - z^2 \langle e^{ib\phi} \rangle^2$ has a finite limit when $\beta = 2$,

$$n_{+-}^{(2)T}(r) \underset{r \rightarrow 0}{=} z^2 r^{-2} + (2\pi z^2)^2 \left[-\frac{1}{2} + C + \ln(\pi z r) \right] + O(r^2), \quad \beta = 2. \quad (5.9)$$

With the known expansion of $K_1(x)$ for small-argument, we verify that the small- r expansion (5.9) is in complete agreement with the exact expression (5.6).

VI. CONCLUSION AND PERSPECTIVES

Using the operator product expansion we have determined the short-distance behavior of the effective potential between two charges Q_1 and Q_2 immersed in a two-dimensional charge-symmetric two-component plasma. We were also able to determine the short-distance behavior of the density profiles around a single external charge Q in the plasma. The coion density profile exhibits a change of behavior at $\beta Q = 2 - \beta$, from a dependence in r^β for $\beta Q < 2 - \beta$ to a dependence in $r^{2-\beta}$ for $\beta Q > 2 - \beta$. On the other hand the leading behavior of the counterion density profile is always $r^{-\beta Q}$. We have also determined the first subleading contribution to the counterion density profile. Specializing to the case $Q = 1$, we obtained the short-distance behavior of the density correlation functions.

This work complements those of Refs. [9, 10, 11] in which the large-distance behavior of the correlations and related quantities were determined. Although there is not (yet) a closed analytical expression for the correlations functions of the two-component plasma, we now have both its large- and short-distance asymptotics.

A possible interesting application of this work is the following. As we mentioned in the introduction, the bulk thermodynamic properties of the two-component plasma have been determined for $\beta < 2$ for a system of point particles [5]. However, Kalinay and Samaj [26] have devised a method to obtain the thermodynamic quantities of the two-component plasma beyond $\beta = 2$ up to $\beta < 3$ for a system where the charges are small disks of radius σ , in the limit $n\sigma^2 \rightarrow 0$ (n is the density). A basic ingredient of their method is the knowledge of the short-distance behavior of the correlation functions in the region $2 < \beta < 4$. Their work is based on the leading behavior, in $r^{-\beta}$, of $n_{+-}(r)$. If the operator product expansion can be used beyond $\beta = 2$, a simple generalization of the analysis of Secs. IV C and V C gives, for

$$2 < \beta < 4,$$

$$n_{++}^{(2)}(r) \underset{r \rightarrow 0}{=} -\frac{4\pi^2 z^4 r^{4-2\beta}}{(\beta-2)^2} \gamma(1 - \frac{\beta}{4})^3 \gamma(-1 + \frac{3\beta}{4}) + O(r^{2-\beta}, r^{8-3\beta}) \quad (6.1)$$

$$n_{+-}^{(2)}(r) \underset{r \rightarrow 0}{=} z^2 r^{-\beta} + z^4 r^{4-2\beta} J(\beta, -\beta, \beta) + O(r^{2-\beta}, r^{8-3\beta}). \quad (6.2)$$

With this expansion, using the method of Ref. [26], the thermodynamic properties of the two-component plasma, in the low-density limit, can be obtained up to $\beta < 10/3$.

Acknowledgments

The author acknowledge partial financial support from COLCIENCIAS grant 1204-05-13625, ECOS-Nord/COLCIENCIAS, and Comité de Investigaciones de la Facultad de Ciencias de la Universidad de los Andes.

APPENDIX A: EXPANSIONS OF $\langle e^{ibQ\phi} \rangle$

In the text we need the expansion of $\langle e^{ibQ\phi} \rangle$ around a regular value of Q and also around the pole $\beta Q = 2$. If $\beta Q < 2$, the expansion is immediate to obtain

$$\langle e^{ib(Q+\epsilon)\phi} \rangle \underset{\epsilon \rightarrow 0}{=} \langle e^{ibQ\phi} \rangle + \epsilon \frac{\partial \langle e^{ibQ\phi} \rangle}{\partial Q} + \frac{\epsilon^2}{2} \frac{\partial^2 \langle e^{ibQ\phi} \rangle}{\partial Q^2} + O(\epsilon^3). \quad (A1)$$

Using the explicit expression (3.3) for $\langle e^{ibQ\phi} \rangle$ we have

$$\begin{aligned} \langle e^{ib(Q+\epsilon)\phi} \rangle \underset{\epsilon \rightarrow 0}{=} \langle e^{ibQ\phi} \rangle & \left\{ 1 + \epsilon \left[I'_b(Q) + \frac{2\beta Q}{4-\beta} \ln \frac{\pi z}{\gamma(b^2)} \right] \right. \\ & \left. + \frac{\epsilon^2}{2} \left[\left(I'_b(Q) + \frac{2\beta Q}{4-\beta} \ln \frac{\pi z}{\gamma(b^2)} \right)^2 + I''_b(Q) + \frac{2\beta}{4-\beta} \ln \frac{\pi z}{\gamma(b^2)} \right] + O(\epsilon^3) \right\} \end{aligned} \quad (A2)$$

where

$$I'_b(Q) = \frac{\partial I_b(Q)}{\partial Q} = b^2 \int_0^\infty \frac{dt}{t} \left[-4Qe^{-2t} + \frac{t \sinh(4b^2 Qt)}{\sinh t \cosh[(1-b^2)t] \sinh(b^2 t)} \right] \quad (A3)$$

and

$$I''_b(Q) = \frac{\partial^2 I_b(Q)}{\partial Q^2} = 4b^2 \int_0^\infty \frac{dt}{t} \left[-e^{-2t} + \frac{b^2 t^2 \cosh(4b^2 Qt)}{\sinh t \cosh[(1-b^2)t] \sinh(b^2 t)} \right]. \quad (A4)$$

If $\beta Q \rightarrow 2$, the expansion of $\langle e^{ibQ\phi} \rangle$ can be done following the same steps of the appendix of Ref. [10]. Let $\beta Q = 2 - \beta\epsilon$. First, we isolate the pole by writing $I_b(Q) = \mathcal{I}_1(\epsilon) + \mathcal{I}_2(\epsilon) + \mathcal{I}_3(\epsilon)$, with

$$\mathcal{I}_1(\epsilon) = \int_0^1 \frac{dt}{t} \left[\frac{\sinh^2[(1 - 2b^2\epsilon)t]}{2 \sinh(b^2t) \sinh t \sinh[(1 - b^2)t]} - 2 \left(\frac{1}{2b^2} - \epsilon \right)^2 b^2 e^{-2t} \right] \quad (\text{A5a})$$

$$\mathcal{I}_2(\epsilon) = \int_1^\infty \frac{dt}{t} \left[\frac{\sinh^2[(1 - 2b^2\epsilon)t]}{2 \sinh(b^2t) \sinh t \sinh[(1 - b^2)t]} - e^{-4b^2\epsilon t} - 2 \left(\frac{1}{2b^2} - \epsilon \right)^2 b^2 e^{-2t} \right] \quad (\text{A5b})$$

$$\mathcal{I}_3(\epsilon) = \int_1^\infty \frac{e^{-4b^2\epsilon t}}{t} dt = \Gamma(0, 4b^2\epsilon) \underset{\epsilon \rightarrow 0}{=} -\ln(4b^2\epsilon) - C + 4b^2\epsilon - 4(b^2\epsilon)^2 + O(\epsilon^3) \quad (\text{A5c})$$

where $\Gamma(\alpha, x) = \int_x^\infty e^{-t} t^{\alpha-1} dt$ is the incomplete Gamma function. The integral $\mathcal{I}_3$ contains the pole at $\epsilon = 0$, while the integrals $\mathcal{I}_1$ and $\mathcal{I}_2$ are regular at $\epsilon = 0$. The expansion of $\mathcal{I}_1$ and $\mathcal{I}_2$ can simply be obtained with their Taylor series around $\epsilon = 0$: $\mathcal{I}_{1,2}(\epsilon) = \mathcal{I}_{1,2}(0) + \epsilon \mathcal{I}'_{1,2}(0) + (\epsilon^2/2) \mathcal{I}''_{1,2}(0) + O(\epsilon^3)$.

Putting together all the contributions from $\mathcal{I}_1$, $\mathcal{I}_2$, $\mathcal{I}_3$ and from the expansion of $(\pi z / \gamma(b^2))^{Q^2\epsilon}$ in (3.3), we finally find

$$\begin{aligned} \langle e^{ibQ\phi} \rangle_{\beta Q \rightarrow 2} &= 2\pi z \langle e^{ib(\frac{1}{2b^2}-1)\phi} \rangle \left\{ \frac{1}{2 - \beta Q} + \frac{R_1(\beta/4)}{\beta} - \frac{4}{\beta(4 - \beta)} \ln \frac{\pi z}{\gamma(\beta/4)} \right. \\ &+ \frac{2 - \beta Q}{\beta^2} \left[R_2(\beta/4) + \frac{\beta}{4 - \beta} \ln \frac{\pi z}{\gamma(\beta/4)} + \frac{1}{2} \left(R_1(\beta/4) - \frac{4}{4 - \beta} \ln \frac{\pi z}{\gamma(\beta/4)} \right)^2 \right] + O((2 - \beta Q)^2) \Big\} \end{aligned} \quad (\text{A6})$$

with

$$R_1(b^2) = 2 \int_0^\infty \frac{dt}{t} \left[e^{-2t} + tb^2 \left(2 - \frac{\cosh t}{\sinh(b^2t) \cosh[(1 - b^2)t]} \right) \right] \quad (\text{A7})$$

$$R_2(b^2) = 2b^2 \int_0^\infty \frac{dt}{t} \left[b^2 t^2 \left(\frac{\cosh t \coth t + \sinh t}{\sinh(b^2t) \cosh[(1 - b^2)t]} - 4 \right) - e^{-2t} \right]. \quad (\text{A8})$$

Actually, in the main text, we only use the expansion up to order $O(1)$, the $O(\beta Q - 2)$ term is given here only for informational purposes.

It is also useful to know the expansion of $\langle e^{ib\phi} \rangle$ when $\beta \rightarrow 2$. From the explicit formula (2.7) we find

$$\langle e^{ib\phi} \rangle_{\beta \rightarrow 2} = \frac{2\pi z}{2 - \beta} - 2\pi z [C + \ln(\pi z)] - \pi z (\beta - 2) [C + \ln(\pi z)] [1 + C + \ln(\pi z)] + O((\beta - 2)^2) \quad (\text{A9})$$

and

$$\langle e^{ib\phi} \rangle_{\beta \rightarrow 2}^2 = (2\pi z)^2 \left[\frac{1}{(\beta-2)^2} + \frac{2}{\beta-2} [C + \ln(\pi z)] + C + \ln(\pi z) + 2[C + \ln(\pi z)]^2 + O(\beta-2) \right]. \quad (\text{A10})$$

In Sec. V C we need the expansion of $I'_b(1)$ when $\beta \rightarrow 2$ which is a pole of first order. To isolate the pole contribution, it is convenient to write $I'_b(1)$ as

$$\frac{4}{\beta} I'_b(1) = \frac{4}{2-\beta} + \int_0^\infty \left[-\frac{4e^{-2t}}{t} + \frac{\sinh(\beta t)}{\sinh t \sinh(\beta t/4) \cosh[(1-\frac{\beta}{4})t]} - 4e^{-(2-\beta)t} \right] dt. \quad (\text{A11})$$

The remaining integral is now convergent when $\beta \rightarrow 2$, and can be easily expanded around $\beta = 2$. We find

$$\frac{4}{\beta} I'_b(1) = \frac{4}{2-\beta} + 4C - 4(\beta-2) \ln 2 + O((\beta-2)^2). \quad (\text{A12})$$

APPENDIX B: COMPUTATION OF $J(\beta Q, -\beta Q, \beta)$

In this appendix we give an explicit expression for the integral

$$J(\beta Q, -\beta Q, \beta) = \int d^2x d^2y \frac{|x|^{\beta Q} |1-y|^{\beta Q}}{|y|^{\beta Q} |1-x|^{\beta Q} |x-y|^\beta}. \quad (\text{B1})$$

In the appendix D of Ref. [20], an integral more general than (B1), with arbitrary powers in each factor is computed. Using the general results from Ref. [20], we find for our particular case that

$$J(\beta Q, -\beta Q, \beta) = s(\beta/2)^2 \{ (J_1^-)^2 + (J_2^-)^2 \} \quad (\text{B2})$$

where

$$J_1^- = -\frac{s(\beta Q/2)}{s(\beta/2)^2} (s(\beta Q/2) J_1^+ + s(\beta(Q+1)/2) J_2^+) \quad (\text{B3})$$

$$J_2^- = -\frac{s(\beta Q/2)}{s(\beta/2)^2} (s(\beta(Q-1)/2) J_1^+ + s(\beta Q/2) J_2^+) \quad (\text{B4})$$

where we used the notation $s(x) = \sin(\pi x)$ and

$$J_1^+ = \frac{\Gamma(1-\frac{\beta}{2})\Gamma(2-\frac{\beta}{2})\Gamma(1-\frac{\beta Q}{2})^2}{\Gamma(2-\frac{\beta(Q+1)}{2})\Gamma(3-\frac{\beta(Q+1)}{2})} \times {}_3F_2 \left(2-\frac{\beta}{2}, -\frac{\beta Q}{2}, 1-\frac{\beta Q}{2}; 2-\frac{\beta(Q+1)}{2}, 3-\frac{\beta(Q+1)}{2}; 1 \right) \quad (\text{B5})$$

$$J_2^+ = \frac{\Gamma(1-\frac{\beta}{2})\Gamma(2-\frac{\beta}{2})\Gamma(1+\frac{\beta Q}{2})^2}{\Gamma(2+\frac{\beta(Q-1)}{2})\Gamma(3+\frac{\beta(Q-1)}{2})} \times {}_3F_2 \left(2-\frac{\beta}{2}, \frac{\beta Q}{2}, 1+\frac{\beta Q}{2}; 2+\frac{\beta(Q-1)}{2}, 3+\frac{\beta(Q-1)}{2}; 1 \right) \quad (\text{B6})$$

where ${}_3F_2(a_1, a_2, a_3; b_1, b_2; z) = \sum_{k=0}^{\infty} \frac{(a_1)_k (a_2)_k (a_3)_k}{(b_1)_k (b_2)_k k!} z^k$ is a generalized hypergeometric function, and $(a)_k = \Gamma(a+k)/\Gamma(a)$ is the Pochhammer symbol.

For the particular case $Q = 1$, the above expressions are simplified:

$$J(\beta, -\beta, \beta) = [s(\beta/2)J_1^+ + s(\beta)J_2^+]^2 + [s(\beta/2)J_2^+]^2 \quad (\text{B7})$$

with

$$J_1^+ = \frac{\Gamma(1 - \frac{\beta}{2})^3 \Gamma(2 - \frac{\beta}{2})}{\Gamma(2 - \beta) \Gamma(3 - \beta)} {}_3F_2\left(1 - \frac{\beta}{2}, 2 - \frac{\beta}{2}, -\frac{\beta}{2}; 2 - \beta, 3 - \beta; 1\right) \quad (\text{B8})$$

$$J_2^+ = \frac{\Gamma(1 - \frac{\beta}{2}) \Gamma(2 - \frac{\beta}{2}) \Gamma(1 + \frac{\beta}{2})^2}{2} {}_3F_2\left(2 - \frac{\beta}{2}, 1 + \frac{\beta}{2}, \frac{\beta}{2}; 2, 3; 1\right). \quad (\text{B9})$$

To study the limit $\beta \rightarrow 2$, we need the expansion of $J(\beta, -\beta, \beta)$ around $\beta = 2$. In the hypergeometric function defining J_2^+ , $\beta = 2$ is a regular point. We can obtain its expansion with the Taylor series

$$\begin{aligned} {}_3F_2\left(2 - \frac{\beta}{2}, 1 + \frac{\beta}{2}, \frac{\beta}{2}; 2, 3; 1\right) &= {}_3F_2(2, 1, 0; 2, 3; 1) \\ &+ (\beta - 2) \left. \frac{\partial {}_3F_2(2 - \frac{\beta}{2}, 1 + \frac{\beta}{2}, \frac{\beta}{2}; 2, 3; 1)}{\partial \beta} \right|_{\beta=2} + O((\beta - 2)^2). \end{aligned} \quad (\text{B10})$$

Deriving under the summation we find

$$\left. \frac{\partial {}_3F_2(2 - \frac{\beta}{2}, 1 + \frac{\beta}{2}, \frac{\beta}{2}; 2, 3; 1)}{\partial \beta} \right|_{\beta=2} = \sum_{k=1}^{\infty} \frac{-1 + C + \psi(2+k)}{(k+1)(k+2)} = \frac{\pi^2}{6} - 1. \quad (\text{B11})$$

Then

$${}_3F_2\left(2 - \frac{\beta}{2}, 1 + \frac{\beta}{2}, \frac{\beta}{2}; 2, 3; 1\right) = 2 + (\beta - 2) \left(\frac{\pi^2}{6} - 1 \right) + O((\beta - 2)^2). \quad (\text{B12})$$

With this result we have the following expansions, needed for the expansion of $J(\beta, -\beta, \beta)$,

$$s(\beta/2)J_2^+ \underset{\beta \rightarrow 2}{=} \pi + \frac{\pi}{12} (6 + \pi^2) (\beta - 2) + O((\beta - 2)^2) \quad (\text{B13})$$

$$s(\beta)J_2^+ \underset{\beta \rightarrow 2}{=} -2\pi - \frac{\pi}{6} (6 + \pi^2) (\beta - 2) + O((\beta - 2)^2) \quad (\text{B14})$$

For the expansion of the hypergeometric function in J_1^+ , it is convenient to write it as

$${}_3F_2\left(1 - \frac{\beta}{2}, 2 - \frac{\beta}{2}, -\frac{\beta}{2}; 2 - \beta, 3 - \beta; 1\right) = 1 - \frac{\frac{\beta}{2}(2 - \frac{\beta}{2})}{2(3 - \beta)} + (1 - \frac{\beta}{2})S(\beta) \quad (\text{B15})$$

with

$$S(\beta) = - \sum_{k=2}^{\infty} \frac{(\beta/2)(2 - \frac{\beta}{2})_{k-2}(2 - \frac{\beta}{2})_{k-1}(2 - \frac{\beta}{2})_k}{2(3 - \beta)_{k-1}(3 - \beta)_k k!}. \quad (\text{B16})$$

This last term can be expanded to the first order using $S(\beta) \underset{\beta \rightarrow 2}{=} S(2) + (\beta - 2)S'(2) + O(\beta - 2)$, with $S'(2) = - \sum_{k=2}^{\infty} \frac{1+C-\psi(k-1)+\psi(k)+\psi(k+1)}{4k(k-1)} = -(1 + \pi^2/12)/2$. Then,

$$S(\beta) \underset{\beta \rightarrow 2}{=} -\frac{1}{2} - \frac{\beta - 2}{2} \left(1 + \frac{\pi^2}{12}\right) + O((\beta - 2)^2) \quad (\text{B17})$$

and finally

$${}_3F_2 \left(1 - \frac{\beta}{2}, 2 - \frac{\beta}{2}, -\frac{\beta}{2}; 2 - \beta, 3 - \beta; 1\right) \underset{\beta \rightarrow 2}{=} \frac{1}{2} - \frac{\beta - 2}{4} + \frac{(\beta - 2)^2}{8} \left(\frac{\pi^2}{6} - 1\right) + O((\beta - 2)^3). \quad (\text{B18})$$

We expanded up to order $(\beta - 2)^2$, since J_1^+ has a pole of order 2 at $\beta = 2$, as opposed to J_2^+ which has a pole of order 1. With this expansion (B18), we find

$$s(\beta/2)J_1^+ \underset{\beta \rightarrow 2}{=} -\frac{2\pi}{\beta - 2} + \pi + \frac{\pi}{6}(3 + \pi^2)(\beta - 2) + O((\beta - 2)^2). \quad (\text{B19})$$

Putting together equations (B13), (B14), and (B19) we finally find

$$J(\beta, -\beta, \beta) \underset{\beta \rightarrow 2}{=} (2\pi)^2 \left[\frac{1}{(\beta - 2)^2} + \frac{1}{\beta - 2} + 1 \right] + O(\beta - 2). \quad (\text{B20})$$

-
- [1] see e.g., S. Samuel, Grand partition function in field theory with applications to sine-Gordon field theory, *Phys. Rev. D* **18**:1916 (1978).
 - [2] A. Zamolodchikov and Al. Zamolodchikov, Factorized S-matrices in two dimensions as the exact solutions of certain relativistic quantum field theory models, *Ann. Phys. (N.Y.)* **120**:253 (1979).
 - [3] C. Destri and H. de Vega, New exact results in affine Toda field theories: Free energy and wave-function renormalizations, *Nucl. Phys. B* **358**:251 (1991).
 - [4] Al. Zamolodchikov, Mass scale in the sine-Gordon model and its reductions, *Int. J. Mod. Phys. A* **10**:1125 (1995).
 - [5] L. Šamaj and I. Travěnek, Thermodynamic properties of the two-dimensional two-component plasma, *J. Stat. Phys.* **101**:713 (2000).
 - [6] S. Lukyanov and A. Zamolodchikov, Exact expectation values of local fields in the quantum sine-Gordon model, *Nucl. Phys. B* **493**:571 (1997).

- [7] S. Lukyanov, Form-factors of exponential fields in the affine $A_{N-1}^{(1)}$ Toda model, *Mod. Phys. Lett.* **12**:2543 (1997).
- [8] S. Lukyanov, Form-factors of exponential fields in the sine-Gordon model, *Phys. Lett. B* **408**:192 (1997).
- [9] L. Šamaj and B. Jancovici, Large-distance behavior of particle correlations in the two-dimensional two-component plasma, *J. Stat. Phys.* **106**:301 (2002).
- [10] L. Šamaj, Anomalous effects of “guest” charges immersed in electrolyte: Exact 2D results, e-print cond-mat/0503201, to be published in J. Stat. Phys. (2005).
- [11] G. Téllez, Guest charges in an electrolyte: renormalized charge, long- and short-distance behavior of the electric potential and density profiles, e-print cond-mat/0504740, to be published in J. Stat. Phys. (2005).
- [12] V. S. Dotsenko and V. A. Fateev, Conformal algebra and multipoint correlation functions in 2D statistical models, *Nucl. Phys. B* **240**:312 (1984).
- [13] V. S. Dotsenko and V. A. Fateev, Four-point correlation functions and the operator algebra in 2D conformal invariant theories with central charge $c \leq 1$, *Nucl. Phys. B* **251**:691 (1985).
- [14] A. Torres and G. Téllez, Finite-size corrections for Coulomb systems in the Debye–Hückel regime, *J. Phys. A: Math. Gen.* **37**:2121 (2004).
- [15] V. Fateev, S. Lukyanov, A. Zamolodchikov and Al. Zamolodchikov, Expectation values of local fields in the Bullough-Dodd model and integrable perturbed conformal field theories, *Nucl. Phys. B* **516**:652 (1998).
- [16] Z. Bajnok, L. Palla, G. Takács and F. Wágner, The k -folded sine-Gordon model in finite volume, *Nucl. Phys. B* **587**:585 (2000).
- [17] W. F. Lu, Sine-Gordon expectation values of exponential fields with variational perturbation theory, *Phys. Lett. B* **602**:261 (2004).
- [18] K. Wilson, Non-Lagrangian models of current algebra, *Phys. Rev.* **179**:1499 (1969).
- [19] V. Fateev, D. Fradkin, S. Lukyanov, A. Zamolodchikov, and Al. Zamolodchikov, Expectation values of descendent fields in the sine-Gordon model, *Nucl. Phys. B* **540**:587 (1999).
- [20] V. Dotsenko, M. Picco, and P. Pujol, Renormalisation-group calculation of correlation functions for the 2D random bond Ising and Potts models, *Nucl. Phys. B* **455**:701 (1995).
- [21] R. Guida and N. Magnoli, Tricritical Ising model near criticality, *Int. J. Mod. Phys. A* **13**:1145 (1998).

- [22] J.-P. Hansen and P. Viot, Two-body correlations and pair formation in the two-dimensional Coulomb gas, *J. Stat. Phys.* **38**:823 (1985).
- [23] F. Cornu and B. Jancovici, On the two-dimensional Coulomb gas, *J. Stat. Phys.* **49**:33 (1987).
- [24] F. Cornu and B. Jancovici, The electrical double layer: A solvable model, *J. Chem. Phys.* **90**:2444 (1989).
- [25] S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products* (Academic, New York, 1965).
- [26] P. Kalinay and L. Šamaj, Thermodynamic Properties of the Two-Dimensional Coulomb Gas in the Low-Density Limit, *J. Stat. Phys.* **106**:857 (2002).